

GRADE V SPONDYLOLISTHESIS ASSOCIATED WITH A PARAORTIC NEUROFIBROMA -A Case report-

ACAROĞLU E., M.D.

YAZICI M., M.D.

SURAT A., M.D.

INTRODUCTION

Spondylolisthesis is defined as a slipping forward of one vertebra on another. Meyerding described a grading system for classifying the percentage of slippage. In this system, Gr I is slippage of 0 to 25 per cent, and Gr IV is slippage greater than 75 per cent. Gr V spondylolisthesis (spondyloptosis) is the most severe form of spondylolisthesis which is mostly seen in L 5. In spondyloptosis, there is vertical descent of the entire vertebral body of the L 5 below the end plate of S 1.

In spondylolisthesis, the indications for surgery primarily include pain and radiculopathy. Additionally in children, the indications for surgery include a 50 per cent slip and the presence of progression and cosmetic problems. The patient with severe spondylolisthesis should have fusion in normal alignment to prevent progression, relieve pain and improve cosmesis.

Case history

A twenty-year old female patient was admitted with the complaints of low-back pain and gait disturbance for six years. She had a history of two attempted fusions without instrumentation. Both failed to give her any relief. Her physical examination revealed tightness of hamstring muscles on both sides and no motor, sensory or sphincteric neurologic findings. SEP examination was found to be normal. Grade V spondylolisthesis was seen on plain x-ray examination. CT and MRI were performed but no underlying pathology was detected.

The patient was decided to be operated on so as to improve her gait and obtain fusion. The first stage of the operation consisted of posterior ileo-sacro-lumbar instrumentation and postero-lateral fusion. Intraoperatively pedicles were seen to be very narrow, transvers diameters being very small. Transpedicular fixation therefore, could not be performed so sublaminar wiring of L3 and L2 and a bi-laminar claw at L1 were used. For the second stage a trans-peritoneal lumbosacral fusion was planned, but as the operation commenced, a firm paraaortic mass extending from the

aortic bifurcation up to the retrogastric area was encountered. The tumor could only be excised partially due to its extension and anterior fusion was performed. The patient's recovery was unevenful for the twelve months follow-up period. Solid osseous union was seen on plain radiographs. She was relieved of all her pain, but had some residual deformity. Histologic examination of the paraortic mass revealed a neurofibroma. retrospectively, the patient was re-evaluated, but any finding suggesting neurofibromatosis was not found. However, pedicle and facet joint changes was assumed to be dystrophic changes seen in neurofibromatosis.

DISCUSSION

Nerofibromatosis is known to be associated with dystrophic changes in the vertebral column and several vertebral deformities. but spondylolisthesis is an unusual spine deformity in neruofibromatosis. This deformity is frequently known to case is the only patient to our knowledge presenting with paraortic neurofibroma and severe spondylolisthesis. Whether the deformity is caused by the tumor or the association is only incidental is very hard to decide. But this case implies that Gr V spondylolisthesis may be associated with neurofibromatosis in children and this possibility should be considered preoperatively.

REFERENCES

- 1 Bradford DS., Boachie-Adjei O.: Treatment of severe spondylolisthesis by anterior and posterior reduction and stabilization. *J. Bone and Joint Surg.*, 72-A: 1060-1066, 1990
- 2 Bradford DS., Lonstein CE.: Moe's textbook of scoliosis and other spinal deformities. Ed. 2, pp. 403-434. Philadelphia, W. B. Saunders, 1987
- 3 Bridwell KH., DeWald RL.: The textbook of spinal surgery. Ed. 1, pp. 457 and 593-604. Philadelphia, J.B. Lippincott, 1991.
- 4 Gaines RW., Nichols WK.: Treatment of spondyloptosis by Cotrel-Dubouset instrumentation. 7th CDI proceedings reports, 123-127, 1990.
- 5 Seitsalo S.: Operative and conservative treatment of moderate spondylolisthesis in young patients. *J. Bone and Joint Surg.*, 72-B: 908-913, 1990.
- 6 Steffee AD., Sitkowski DJ.: Reduction and stabilization of grade IV spondylolisthesis. *Clin. Orthop.*, 227: 82-89, 1988.

HIGH THORACIC CONGENITAL KYPHOSIS ASSOCIATED WITH A HIGH MEDIASTINAL NEUROENTERIC CYST (A Case Report)

SURAT A., M.D, ACAROĞLU E., M.D, YAZICI M., M.D, LEBLEBİCİOĞLU G., M.D.

A case of high thoracic kyphosis, that was found to be associated with a neuroenteric cyst located at the posterior mediastinum is presented.

Total excision of the cystic lesion immediately anterior to the vertebral column could be accomplished by a right thoracotomy and anterior as well as posterior in-situ fusions were performed for the rapidly progressing deformity.

Congenital kyphotic deformities of the spinal column located at the high thoracic area are known to be rapidly progressing lesions with dismal prognoses. The natural history of these deformities are often associated with a high rate of spinal compression and neurologic involvement. Posterior fusions alone are sufficient for neither halting the progression of the deformities, nor decompression of the spinal cord that is usually pressed to the posterior sides of the vertebral bodies. Hence anterior and posterior fusions should be the treatment of choice and should be contemplated as early as possible.

Neuroenteric cysts are a subgroup of split notocord syndromes, formed by the adhesions between the neural tube and the gut. They may be in continuity with the gut or the subarachnoid space or rarely both, and may as well form an enteric fistula. The most common locations are the posterior mediastinum and the retroperitoneal region.

Case Report:

A two and one half year old girl was brought to the our department with her parents observation of a rapidly progressing hump at the base of her neck. The deformity was noticed three months ago. The patient had begun walking by the age of thirteen months but was not fully continent as yet.

She had a small hump at her high thoracic area, no neurologic defect could be detected. Radiologic examination revealed a high thoracic konjenital kyphosis measuring 57 degrees from T1 to T5 and lower thoracic hypo-kyphosis. Suspecting of a dysraphic spine Magnetic Resonans Imaging (MRI) of the region and a

control visit three months later were scheduled.

MRI revealed a huge cystic mass located at the posterior mediastinum in the vicinity of the vertebral deformity that was thought to be a neuroenteric cyst and the patient's deformity has progressed to 68 degrees during the three months. Neither myelographic examination nor the three months. Neither myelographic examination nor the upper GI series demonstrated any leakage of dye into the cystic cavity.

The decision to operate was mainly based on the rapid progression of the vertebral deformity. A right thoracotomy through the third right rib, elevating the scapula, revealed a retropleural cystic mass filled with a clear and very viscous fluid. The cyst could be thoroughly excised and an anterior fusion using the excised rib as a strut graft was performed after discectomies. Her post-operative recovery period was eventless and a posterior fusion was performed four weeks later. The patient was discharged from the hospital wearing a Minerva type cast. The cast was removed at three months follow-up visit and despite some displacement of the anterior strut graft a bony fusion was achieved.

Histologic examination of the mass revealed a cyst cavity lined up with primitive neural like elements and the diagnosis of a neuroenteric cyst was confirmed.

DISCUSSION

Congenital kyphosis is a well known clinical entity and if located at the high thoracic region carries a very high risk of cord compression and neurologic involvement. Results obtained by braces have been very unsatisfactory and early surgical fusion of the deformity is usually necessary. Posterior approaches per-se have very high rates of pseudoarthroses and are nearly always insufficient for spinal decompression. Both ante-

rior and posterior fusions and if necessary anterior decompression should be the treatment of choice.

Congenital deformities of the vertebral column are frequently associated with dysraphic changes of the spinal cord, so MRI examination has long been a standard preoperative procedure. As can be seen from our case it has proven helpful in delineating other possible abnormalities, that may prove very difficult to deal with if not prepared.

Whether the association of the cyst and the vertebral congenital deformity is purely incidental or if there is a cause-effect relation between the two is very hard to decide. Neuroenteric cysts are thought to be caused by adhesions between the neural tube and the endoderm that will form the fore-gut and these adhesions might probably have caused formation defects at the anterior portions of the somites.

REFERENCES:

1. Beardmore H.E, Wigglesworth F.W.; Vertebral anomalies and alimentary duplications. *Ped. Clin. North Am.*, 5, 547- 474, 1958.
2. Bentley F.F, Smith J.R.; Developmental posterior enteric remnants and spinal malformations, The split notochord syndrome. *Arch. Dis. Child.*, 35, 76- 86, 1960.
3. Ehrenhaft J.L.; The development of vertebral column as related to certain congenital and pathological changes. *Surg. Gynecol. Obstet*, 76, 202-92, 1943.
4. Fallon M, Gordon A.R.G, Lendrum A.C.; Mediastinal cysts of foregut origin associated with vertebral anomalies. *Br. J. Surg*, 41, 520-33, 1953.
5. O'Connor J.F, Cranley W.R, McCarten K.M. et al.; Radiographic manifestations of congenital anomalies of the spine. *Radiol. Clin. North Am.*, 29, 407-29, 1991.